


## Control method: Cage trapping of feral cats (with transport) followed by killing

Assumptions:	<ul style="list-style-type: none"> <li>▪ Best practice is followed in accordance with CAT002.</li> <li>▪ Traps are set in the evening and checked in the morning.</li> <li>▪ Efforts are made to locate and kill any kittens if lactating queen is caught.</li> <li>▪ Shooting is the usual method of euthanasia although lethal injection is sometimes used. In some situations it is necessary to transport the trapped animals to another location for euthanasia.</li> <li>▪ Cage trapping is often used when cats are at high densities around human settlements e.g. around rubbish tips, camping grounds and less often when cat densities are low in areas away from human habitation.</li> </ul>
--------------	---

### PART A: assessment of overall welfare impact



<b>SCORE FOR PART A:</b>	<b>5</b>
Summary of evidence:	Note that the cat will spend additional time in the trap compared with being shot at the site of capture but this is still likely to be less than 24 hours.
Domain 1	Traps are set in the evening and checked in the morning. Food bait is provided but no water.
Domain 2	Assumes traps are not set in bad weather and are placed in shaded areas.
Domain 3	There is the potential for minor injuries to be sustained, usually self-inflicted abrasions to the face <sup>1,2</sup> .
Domain 4	There will be some restraint stress but cats quickly recover from this if released. The physiological response to capture has been found to be lower in animals caught in cage traps compared with leg-hold traps <sup>3</sup> . In foxes, cage traps caused an increase in cortisol compared with animals that were not trapped but this was lower than individuals caught in leg-hold traps <sup>4</sup> . There will be some exertion from struggling within the trap; however this will be lower compared with animals held by leg-hold traps <sup>4</sup> . Long entrapment periods could result in disruption of natural behaviour and motivational systems <sup>5</sup> . Although the cage will be covered, transporting the cat to another site will cause additional stress
Domain 5	It is likely that the animal will experience an elevation in anxiety and distress during trapping and transportation <sup>6</sup> . Evidence that animals can be recaptured in cage traps may indicate that overall the impact of trapping alone is not high or long-term <sup>2</sup> .

### PART B: assessment of mode of death – shooting (head shot)

Time to insensibility (minus any lag time)				
Immediate to seconds	Minutes	Hours	Days	Weeks
Level of suffering (after application of the method that causes death but before insensibility)				
No suffering	Mild suffering	Moderate suffering	Severe suffering	Extreme suffering

### PART B: assessment of mode of death – lethal injection

Time to insensibility (minus any lag time)				
Immediate to seconds	Minutes	Hours	Days	Weeks
Level of suffering (after application of the method that causes death but before insensibility)				
No suffering	Mild suffering	Moderate suffering	Severe suffering	Extreme suffering

<b>SCORE FOR PART B:</b>	<b>Shooting (head shot) - B</b>
Summary of evidence:	
Duration –	With head shots, a properly placed shot will result in immediate insensibility <sup>7,8,9</sup> .

Suffering –	The approach of a human to a trapped cat will cause some distress <sup>10</sup> . A well-placed head shot which causes immediate insensibility should not cause any additional suffering.
-------------	---

<b>SCORE FOR PART B:</b>	<b>Lethal injection - D</b>
Summary of evidence:	
Duration –	The duration will start from approach of human followed by an intramuscular injection (IM) of sedative and/or anaesthetic agent administered with the aid of a crush/squeeze cage or with a pole syringe. Heavy sedation/loss of consciousness occurs approx. 15 minutes afterwards.
Suffering –	The approach of a human to a trapped cat will cause some distress <sup>10</sup> . Also there will be some pain associated with the IM injection. The animal is then not approached again until fully sedated or unconscious. An overdose of barbiturate administered by the intravenous, intraperitoneal or intracardiac routes should cause no suffering in an anaesthetised or heavily sedated cat <sup>8</sup> .

## Summary

<b>CONTROL METHOD:</b>	<b>Cage trapping of feral cats (with transport) followed by killing</b>
<b>OVERALL HUMANENESS SCORE:</b>	<b>Cage trapping with shooting (head shot) – 5B Cage trapping with lethal injection – 5D</b>

## Bibliography

- Short, J., Turner, B. & Risbey, D. (2002). Control of feral cats for nature conservation. III. Trapping. *Wildlife Research* **29**, 475-487
- Molsher, R.L. (2001). Trapping and demographics of feral cats (*Felis catus*) in central New South Wales. *Wildlife Research* **28**, 631-636
- Iossa, G., Soulsbury, C.D. & Harris, S. (2007). Mammal trapping: a review of animal welfare standards of killing and restraining traps. *Animal Welfare* **16**, 335-352
- White, P.J., Kreeger, T.J., Seal, U.S. & Tester, J.R. (1991). Pathological responses of red foxes to capture in box traps. *Journal of Wildlife Management* **55**, 75-80
- Schutz, K.E. et al. (2006). Behavioural and physiological responses of trap-induced stress in European badgers. *Journal of Wildlife Management* **70**, 884-891
- Williams, L.S., Levy, J.K., Robertson, S.A., Cistola, A.M. & Centonze, L.A. (2002). Use of the anesthetic combination of tiletamine, zolazepam, ketamine, and xylazine for neutering feral cats. *Journal of the American Veterinary Medical Association* **220**, 1491-1495
- Gregory, N. (2004). *Physiology and behaviour of animal suffering*. (Blackwell: Oxford, UK).
- Association, A.V.M. (2001). 2000 Report of the AVMA Panel on Euthanasia. *Journal of the American Veterinary Medical Association* **218**, 669-696
- Longair, J. et al. (1991). Guidelines for euthanasia of domestic animals by firearms. *Canadian Veterinary Journal* **32**, 724-726
- Bakken, M., Moe, R.O., Smith, A.J. & Selle, G.M.E. (1999). Effects of environmental stressors on deep body temperature and activity levels in silver fox vixens (*Vulpes vulpes*). *Applied Animal Behaviour Science* **64**, 141-151